



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ation, and weight will descend with an acceleration $a-b$ times the angular acceleration.

99. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

In a triangle ABC , base $=b$, area $=\Delta$, the principal moments of inertia at the centroid are $\frac{1}{72}m[a^2+b^2+c^2 \pm \sqrt{(a^4+b^4+c^4-a^2b^2-a^2c^2-b^2c^2)}]$ and the principal axes at this point make with the base AC an angle θ given by

$$\tan 2\theta = \frac{4(c^2 - a^2)\Delta}{(a^2 - c^2)^2 - b^2(a^2 + c^2) + 2b^4}.$$

Solution by the PROPOSER.

Let O be the centroid and transform from the rectangular axes Ox, Oy , to the oblique axes Ox, OB .

Also let $(b/6AD)(2AD-3y)=x'$.

$$\begin{aligned}\text{Then } \Sigma mxy &= \rho \sin^2 D \int_{-\frac{1}{3}AD}^{\frac{2}{3}AD} \int_{-x'}^{x'} (x+y\cos D) y dy dx = \frac{1}{3} b \rho AD^3 \sin^2 D \cos D \\ &= \frac{1}{18} m AD^2 \sin D \cos D.\end{aligned}$$

$$\Sigma mx^2 = \rho \sin D \int_{-\frac{1}{3}AD}^{\frac{2}{3}AD} \int_{-x'}^{x'} (x+y\cos D)^2 dy dx = \frac{1}{2} m (3b^2 + 4AD^2 \cos^2 D).$$

$$\Sigma my^2 = \rho \sin^3 D \int_{-\frac{1}{3}AD}^{\frac{2}{3}AD} \int_{-x'}^{x'} y^2 dy dx = \frac{1}{8} m AD^2 \sin^2 D.$$

$$\therefore \tan 2\theta = \frac{4AD^2 \sin 2D}{3b^2 + 4AD^2 \cos 2D}.$$


$$\text{But } \sin D = a \sin C / AD, \cos D = \frac{4AD^2 + b^2 - 4a^2}{4b \cdot AD}.$$

$$\sin 2D = \frac{a \sin C (4AD^2 + b^2 - 4a^2)}{2b \cdot AD^2} = \frac{2(c^2 - a^2)\Delta}{b^2 \cdot AD^2}.$$

$$\cos 2D = \frac{AD^2 - 2a^2 \sin^2 C}{AD^2} = \frac{2(a^2 - c^2) - 2b^2(a^2 + c^2) + b^4}{4b^2 \cdot AD^2}.$$

$$\therefore \tan 2\theta = \frac{4(c^2 - a^2)\Delta}{(a^2 - c^2)^2 - b^2(a^2 + c^2) + 2b^4}.$$

$$A \cos^2 \theta + B \sin^2 \theta = \frac{1}{2} m (3b^2 + 4AD^2 \cos^2 D).$$



$$A \sin^2 \theta + B \cos^2 \theta = \frac{1}{18} m A D^2 \sin^2 D.$$

$$\therefore A + B = \frac{1}{72} m (3b^2 + 4AD^2).$$

$$(A - B) \cos 2\theta = \frac{1}{72} m (3b^2 + 4AD^2 \cos 2D).$$

$$\therefore A = \frac{1}{72} m [a^2 + b^2 + c^2 + 2\sqrt{(a^4 + b^4 + c^4 - a^2 b^2 - a^2 c^2 - b^2 c^2)}].$$

$$B = \frac{1}{72} m [a^2 + b^2 + c^2 - 2\sqrt{(a^4 + b^4 + c^4 - a^2 b^2 - a^2 c^2 - b^2 c^2)}].$$

MISCELLANEOUS.

85. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Prove that at least one of the three sides of a rational right triangle must be divisible by 5.

II. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.; HON. JOSIAH H. DRUMMOND, LL. D., Portland, Me.; and the PROPOSER.

Rational right triangles are divided into three kinds: (1), prime integral; (2), composite integral; and (3), fractional.

It is evident that prime integral right triangles are the basis of work.

To produce a prime integral right triangle we take *two integers prime to each other, one odd and the other even*. Then, twice their product will give one of the legs, the difference of their squares will give the other of the legs, and the sum of their squares will be the hypotenuse; or taking p and q as the two integers, the respective sides are $2pq$, $p^2 - q^2$, and $p^2 + q^2$.

This process will produce any prime integral right triangle.

The squares of even numbers end in 4, 6, and 0, and of odd numbers in 1, 9, and 5.

When the two prime integers, or p and q end, respectively, in 0 and an odd figure, or in 5 and an even figure, their product, or pq of the side $2pq$ contains the factor 5; the other sides being odd numbers *not* ending in 5.

When the squares of the two prime integers, or p^2 and q^2 end, respectively, in 1 and 6 or in 9 and 4, the difference of these squares, or the side p^2 and q^2 ends in 5, and is, therefore, divisible by 5.

When the squares of the two prime integers end, respectively, in 1 and 4 or in 9 and 6, the sum of these squares, or the side $p^2 + q^2$ ends in 5.

The above contains all the combinations of any two prime integers, one odd and the other even, according to the formation of the sides of prime integral right triangles.

\therefore In every *prime* integral right triangle one of the sides and *only one* is divisible by 5.

In composite integral and fractional right triangles either *one* or *all* of the sides must be divisible by 5. For, in order to have more than one side divisible by 5, the highest common factor of the three sides must contain the factor 5.

\therefore In rational right triangles either *one* or *all* of the sides will be divisible by 5.